

# Recent BES Results on Hadron Spectroscopy

Chang-Zheng Yuan (for the BES Collaboration)<sup>a\*</sup>

<sup>a</sup>Institute of High Energy Physics, Chinese Academy of Sciences,  
P.O. Box 918-1, Beijing 100049, China

We present recent results from the BES experiment on the observation of the  $Y(2175)$  in  $J/\psi \rightarrow \phi f_0(980)\eta$ , study of  $\eta(2225)$  in  $J/\psi \rightarrow \gamma\phi\phi$ , and the production of  $X(1440)$  recoiling against an  $\omega$  or a  $\phi$  in  $J/\psi$  hadronic decays. The observation of  $\psi(2S)$  radiative decays is also presented.

## 1. Introduction

The analyses reported in this talk were performed using either a sample of  $58 \times 10^6$   $J/\psi$  events or a sample of  $14 \times 10^6$   $\psi(2S)$  events collected with the upgraded Beijing Spectrometer (BESII) detector [1] at the Beijing Electron-Positron Collider (BEPC).

## 2. The $Y(2175)$ in $J/\psi \rightarrow \phi f_0(980)\eta$ [2]

A new structure, denoted as  $Y(2175)$  and with mass  $m = 2.175 \pm 0.010 \pm 0.015$  GeV/ $c^2$  and width  $\Gamma = 58 \pm 16 \pm 20$  MeV/ $c^2$ , was observed by the BaBar experiment in the  $e^+e^- \rightarrow \gamma_{ISR}\phi f_0(980)$  initial-state radiation process [3,4]. This observation stimulated some theoretical speculation that this  $J^{PC} = 1^{--}$  state may be an  $s$ -quark version of the  $Y(4260)$  since both of them are produced in  $e^+e^-$  annihilation and exhibit similar decay patterns [5,6].

Here we report the observation of the  $Y(2175)$  in the decays of  $J/\psi \rightarrow \eta\phi f_0(980)$ , with  $\eta \rightarrow \gamma\gamma$ ,  $\phi \rightarrow K^+K^-$ ,  $f_0(980) \rightarrow \pi^+\pi^-$ . A four-constraint energy-momentum conservation kinematic fit is performed to the  $K^+K^-\pi^+\pi^-\gamma\gamma$  hypothesis for the selected four charged tracks and two photons.  $\eta \rightarrow \gamma\gamma$  candidates are defined as  $\gamma$ -pairs with  $|M_{\gamma\gamma} - 0.547| < 0.037$  GeV/ $c^2$ , a  $\phi$  signal

is defined as  $|m_{K^+K^-} - 1.02| < 0.019$  GeV/ $c^2$ , and in the  $\pi^+\pi^-$  invariant mass spectrum, candidate  $f_0(980)$  mesons are defined by  $|m_{\pi^+\pi^-} - 0.980| < 0.060$  GeV/ $c^2$ . The  $\phi f_0(980)$  invariant mass spectrum for the selected events is shown in Fig. 1, where a clear enhancement is seen around 2.18 GeV/ $c^2$ . Fit with a Breit-Wigner and a polynomial background yields  $52 \pm 12$  signal events and the statistical significance is found to be  $5.5\sigma$  for the signal. The mass of the structure is determined to be  $M = 2.186 \pm 0.010$  (stat)  $\pm 0.006$  (syst) GeV/ $c^2$ , the width is  $\Gamma = 0.065 \pm 0.023$  (stat)  $\pm 0.017$  (syst) GeV/ $c^2$ , and the product branching ratio is  $\mathcal{B}(J/\psi \rightarrow \eta Y(2175)) \cdot \mathcal{B}(Y(2175) \rightarrow \phi f_0(980)) \cdot \mathcal{B}(f_0(980) \rightarrow \pi^+\pi^-) = (3.23 \pm 0.75$  (stat)  $\pm 0.73$  (syst))  $\times 10^{-4}$ . The mass and width are consistent with BaBar's results.

## 3. The $\eta(2225)$ in $J/\psi \rightarrow \gamma\phi\phi$ [7]

Structures in the  $\phi\phi$  invariant-mass spectrum have been observed by several experiments both in the reaction  $\pi^-p \rightarrow \phi\phi n$  [8] and in radiative  $J/\psi$  decays [9–11]. The  $\eta(2225)$  was first observed by the MARK-III collaboration in  $J/\psi$  radiative decays  $J/\psi \rightarrow \gamma\phi\phi$ . A fit to the  $\phi\phi$  invariant-mass spectrum gave a mass of 2.22 GeV/ $c^2$  and a width of 150 MeV/ $c^2$  [9]. An angular analysis of the structure found it to be consistent with a  $0^{-+}$  assignment. It was subsequently observed by the DM2 collaboration, also in  $J/\psi \rightarrow \gamma\phi\phi$  decays [10,11].

We present results from a high statistics study

\*Invited talk at the International workshop on  $e^+e^-$  collisions from Phi to Psi, Laboratori Nazionali di Frascati, Italy, 7-10 April, 2008. Supported in part by the 100 Talents Program of CAS under Contract No. U-25 and by National Natural Science Foundation of China under Contract No. 10491303.




Figure 1. The  $\phi f_0(980)$  invariant mass distribution of the data (points with error bars) and the fit (solid curve) with a Breit-Wigner function and polynomial background; the dashed curve indicates the background function.

of  $J/\psi \rightarrow \gamma\phi\phi$  in the  $\gamma K^+ K^- K_S^0 K_L^0$  final state, with the  $K_L^0$  missing and reconstructed with a one-constraint kinematic fit. After kinematic fit, we require both the  $K^+ K^-$  and  $K_S^0 K_L^0$  invariant masses lie within the  $\phi$  mass region ( $|M(K^+ K^-) - m_\phi| < 12.5 \text{ MeV}/c^2$  and  $|M(K_S^0 K_L^0) - m_\phi| < 25 \text{ MeV}/c^2$ ). The  $\phi\phi$  invariant mass distribution is shown in Fig. 2. There are a total of 508 events with a prominent structure around  $2.24 \text{ GeV}/c^2$ .

A partial wave analysis of the events with  $M(\phi\phi) < 2.7 \text{ GeV}/c^2$  was performed. The two-body decay amplitudes in the sequential decay process  $J/\psi \rightarrow \gamma X$ ,  $X \rightarrow \phi\phi$ ,  $\phi \rightarrow K^+ K^-$  and  $\phi \rightarrow K_S^0 K_L^0$  are constructed using the covariant helicity coupling amplitude method. The intermediate resonance  $X$  is described with the normal Breit-Wigner propagator  $BW = 1/(M^2 - s - iMT)$ , where  $s$  is the  $\phi\phi$  invariant mass-squared and  $M$  and  $\Gamma$  are the resonance's mass and width. When  $J/\psi \rightarrow \gamma X$ ,  $X \rightarrow \phi\phi$  is fitted with both the  $\phi\phi$  and  $\gamma X$  systems in a  $P$ -wave, which corresponds to a pseudoscalar  $X$  state, the fit gives  $196 \pm 19$  events with mass  $M = 2.24^{+0.03+0.03}_{-0.02-0.02} \text{ GeV}/c^2$ , width  $\Gamma = 0.19 \pm$




Figure 2. The  $K^+ K^- K_S^0 K_L^0$  invariant mass distribution for  $J/\psi \rightarrow \gamma\phi\phi$  candidate events. The dashed histogram is the phase space invariant mass distribution, and the dotted curve indicates how the acceptance varies with the  $\phi\phi$  invariant mass.

$0.03^{+0.04}_{-0.06} \text{ GeV}/c^2$ , and a statistical significance larger than  $10\sigma$ , and a product branching fraction of:  $\mathcal{B}(J/\psi \rightarrow \gamma\eta(2225)) \cdot \mathcal{B}(\eta(2225) \rightarrow \phi\phi) = (4.4 \pm 0.4 \pm 0.8) \times 10^{-4}$ .

The presence of a signal around  $2.24 \text{ GeV}/c^2$  and its pseudoscalar character are confirmed, and the mass, width, and branching fraction are in good agreement with previous experiments.

#### 4. The $X(1440)$ in $J/\psi$ hadronic decays [12]

A pseudoscalar gluonium candidate, the so-called  $E/\iota(1440)$ , was observed in  $p\bar{p}$  annihilation in 1967 [13] and in  $J/\psi$  radiative decays in the 1980's [14–16]. The study of the decays  $J/\psi \rightarrow \{\omega, \phi\} K\bar{K}\pi$  is a useful tool in the investigation of quark and possible gluonium content of the states around  $1.44 \text{ GeV}/c^2$ . Here we investigate the possible structure in the  $K\bar{K}\pi$  final state in  $J/\psi$  hadronic decays at around  $1.44 \text{ GeV}/c^2$ .

In this analysis,  $\omega$  mesons are observed in the  $\omega \rightarrow \pi^+ \pi^- \pi^0$  decay,  $\phi$  mesons in the  $\phi \rightarrow K^+ K^-$  decay, and other mesons are detected in the decays:  $K_S^0 \rightarrow \pi^+ \pi^-$ ,  $\pi^0 \rightarrow \gamma\gamma$ .  $K\bar{K}\pi$  could be

$K_S^0 K^\pm \pi^\mp$  or  $K^+ K^- \pi^0$ .

Figures 3 and 4 show the  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  invariant mass spectra after  $\omega$  selection ( $|m_{\pi^+ \pi^- \gamma \gamma} - m_\omega| < 0.04 \text{ GeV}/c^2$ ) or  $\phi$  signal selection ( $|m_{K^+ K^-} - m_\phi| < 0.015 \text{ GeV}/c^2$ ). Clear  $X(1440)$  signal is observed recoiling against the  $\omega$ , and there is no significant signal recoiling against a  $\phi$ .

The  $K_S^0 K^\pm \pi^\mp$  invariant mass distribution in  $J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$  (Fig. 3(b)) is fitted with a BW function convoluted with a Gaussian mass resolution function ( $\sigma = 7.44 \text{ MeV}/c^2$ ) to represent the  $X(1440)$  signal and a third-order polynomial background function. The mass and width obtained from the fit are  $M = 1437.6 \pm 3.2 \text{ MeV}/c^2$  and  $\Gamma = 48.9 \pm 9.0 \text{ MeV}/c^2$ , and the fit yields  $249 \pm 35$  events. Using the efficiency of 1.45% determined from a uniform phase space MC simulation, we obtain the branching fraction to be  $\mathcal{B}(J/\psi \rightarrow \omega X(1440)) \cdot \mathcal{B}(X(1440) \rightarrow K_S^0 K^+ \pi^- + c.c.) = (4.86 \pm 0.69 \pm 0.81) \times 10^{-4}$ , where the first error is statistical and the second one systematic.

For  $J/\psi \rightarrow \omega K^+ K^- \pi^0$  mode, by fitting the  $K^+ K^- \pi^0$  mass spectrum in Fig. 3(c) with same functions, we obtain the mass and width of  $M = 1445.9 \pm 5.7 \text{ MeV}/c^2$  and  $\Gamma = 34.2 \pm 18.5 \text{ MeV}/c^2$ , and the number of events from the fit is  $62 \pm 18$ . The efficiency is determined to be 0.64% from a phase space MC simulation, and the branching fraction is  $\mathcal{B}(J/\psi \rightarrow \omega X(1440)) \cdot \mathcal{B}(X(1440) \rightarrow K^+ K^- \pi^0) = (1.92 \pm 0.57 \pm 0.38) \times 10^{-4}$ , in good agreement with the isospin symmetry expectation from  $J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$  mode.

The distribution of  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  invariant mass spectra recoiling against the  $\phi$  signal are shown in Fig. 4, and there is no evidence for  $X(1440)$ . The upper limits on the branching fractions at the 90% C.L. are  $\mathcal{B}(J/\psi \rightarrow \phi X(1440) \rightarrow \phi K_S^0 K^+ \pi^- + c.c.) < 1.93 \times 10^{-5}$  and  $\mathcal{B}(J/\psi \rightarrow \phi X(1440) \rightarrow \phi K^+ K^- \pi^0) < 1.71 \times 10^{-5}$ .

In conclusion, the mass and width of the  $X(1440)$  are measured, which are in agreement with previous measurements; the branching fractions we measured are also in agreement with the DM2 and MARK-III results. The significant signal in  $J/\psi \rightarrow \omega K \bar{K} \pi$  mode and the missing signal in  $J/\psi \rightarrow \phi X$  mode may indicate the  $s\bar{s}$  compo-

nent in the  $X(1440)$  is not significant.

## 5. $\psi(2S)$ radiative decays

Besides conventional meson and baryon states, QCD also predicts a rich spectrum of glueballs, hybrids, and multi-quark states in the 1.0 to 2.5  $\text{GeV}/c^2$  mass region. Therefore, searches for the evidence of these exotic states play an important role in testing QCD. The radiative decays of  $\psi(2S)$  to hadrons are expected to contribute about 1% to the total  $\psi(2S)$  decay width [17]. However, the measured channels only sum up to about 0.05% [18].

We measured the decays of  $\psi(2S)$  into  $\gamma p \bar{p}$ ,  $\gamma 2(\pi^+ \pi^-)$ ,  $\gamma K_S^0 K^+ \pi^- + c.c.$ ,  $\gamma K^+ K^- \pi^+ \pi^-$ ,  $\gamma K^{*0} K^- \pi^+ + c.c.$ ,  $\gamma K^{*0} \bar{K}^{*0}$ ,  $\gamma \pi^+ \pi^- p \bar{p}$ ,  $\gamma 2(K^+ K^-)$ ,  $\gamma 3(\pi^+ \pi^-)$ , and  $\gamma 2(\pi^+ \pi^-) K^+ K^-$ , with the invariant mass of the hadrons ( $m_{hs}$ ) less than 2.9  $\text{GeV}/c^2$  for each decay mode [19]. The differential branching fractions are shown in Fig. 5. The branching fractions below  $m_{hs} < 2.9 \text{ GeV}/c^2$  are given in Table 1, which sum up to 0.26% of the total  $\psi(2S)$  decay width. We also analyzed  $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$  and  $\gamma K^+ K^-$  modes to study the resonances in  $\pi^+ \pi^-$  and  $K^+ K^-$  invariant mass spectrum. Significant signals for  $f_2(1270)$  and  $f_0(1710)$  were observed, but the low statistics prevent us from drawing solid conclusion on the other resonances [20].

## 6. Summary

Using the 58 M  $J/\psi$  and 14 M  $\psi(2S)$  events samples taken with the BESII detector at the BEPC storage ring, BES experiment provided many interesting results in charmonium decays, including the observation of the  $Y(2175)$ ,  $\eta(2225)$ ,  $X(1440)$ , and many  $\psi(2S)$  radiative decays. These results shed light on the understanding of strong interaction sector of the Standard Model.

## REFERENCES

1. J. Z. Bai *et al.* [BES Collaboration], Nucl. Instr. Meth. A **458**, 627 (2001).
2. M. Ablikim *et al.* [BES Collaboration], Phys. Rev. Lett. **100**, 102003 (2008).




Figure 3. The  $K\bar{K}\pi$  invariant mass distribution for  $J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$  (b) and  $\omega K^+ K^- \pi^0$  (c) candidate events. The curves are the best fit.




Figure 4. The  $K_S^0 K^\pm \pi^\mp$  (left) and  $K^+ K^- \pi^0$  (right) invariant mass recoiling against the  $\phi$  in  $J/\psi \rightarrow \phi K\bar{K}\pi$  mode.




Figure 5. Differential branching fractions for  $\psi(2S) \rightarrow \gamma p\bar{p}$ ,  $\gamma 2(\pi^+\pi^-)$ ,  $\gamma K^+K^-\pi^+\pi^-$ , and  $\gamma K_S^0 K^+\pi^- + c.c.$ . Here  $m_{hs}$  is the invariant mass of the hadrons. For each point, the smaller longitudinal error is the statistical error, while the bigger one is the total error.

Table 1

Branching fractions for  $\psi(2S) \rightarrow \gamma + \text{hadrons}$  with  $m_{hs} < 2.9 \text{ GeV}/c^2$ , where the upper limits are determined at the 90% C.L.

| Mode                            | $\mathcal{B}(\times 10^{-5})$ |
|---------------------------------|-------------------------------|
| $\gamma p\bar{p}$               | $2.9 \pm 0.4 \pm 0.4$         |
| $\gamma 2(\pi^+\pi^-)$          | $39.6 \pm 2.8 \pm 5.0$        |
| $\gamma K_S^0 K^+\pi^- + c.c.$  | $25.6 \pm 3.6 \pm 3.6$        |
| $\gamma K^+K^-\pi^+\pi^-$       | $19.1 \pm 2.7 \pm 4.3$        |
| $\gamma K^{*0} K^+\pi^- + c.c.$ | $37.0 \pm 6.1 \pm 7.2$        |
| $\gamma K^{*0} \bar{K}^{*0}$    | $24.0 \pm 4.5 \pm 5.0$        |
| $\gamma \pi^+\pi^- p\bar{p}$    | $2.8 \pm 1.2 \pm 0.7$         |
| $\gamma K^+K^- K^+K^-$          | $< 4$                         |
| $\gamma 3(\pi^+\pi^-)$          | $< 17$                        |
| $\gamma 2(\pi^+\pi^-) K^+K^-$   | $< 22$                        |

3. B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **74**, 091103 (2006).
4. B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **76**, 012008 (2007).
5. B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. Lett. **95**, 142001 (2005).
6. C. Z. Yuan *et al.* [Belle Collaboration], Phys. Rev. Lett. **99**, 182004 (2007).
7. M. Ablikim *et al.* [BES Collaboration], Phys. Lett. B **662**, 330 (2008).
8. A. Etkin *et al.*, Phys. Lett. B **201**, 568 (1988).
9. Z. Bai *et al.* [MARK-III Collaboration], Phys. Rev. Lett. **65**, 1309 (1990).
10. D. Bisello *et al.* [DM2 Collaboration], Phys. Lett. B **179**, 294 (1986).
11. D. Bisello *et al.* [DM2 Collaboration], Phys. Lett. B **241**, 617 (1990).
12. M. Ablikim *et al.* [BES Collaboration], Phys. Rev. D **77**, 032005 (2008).
13. P. H. Baillon *et al.*, NC 50A, 393 (1967).
14. D. L. Scharre *et al.*, Phys. Lett. B **97**, 329 (1980).
15. C. Edwards *et al.*, Phys. Rev. Lett. **49**, 259 (1982); Phys. Rev. Lett. **50**, 219 (1982).
16. J. E. Augustin *et al.* [DM2 Collaboration], Phys. Rev. D **42**, 10 (1990).
17. P. Wang, C. Z. Yuan, and X. H. Mo, Phys. Rev. D **70**, 114014 (2004).
18. W.-M. Yao *et al.* [Particle Data Group], J. Phys. G **33**, 1 (2006).
19. M. Ablikim *et al.* [BES Collaboration], Phys. Rev. Lett. **99**, 011802 (2007).
20. M. Ablikim *et al.* [BES Collaboration], arXiv:0710.2324 [hep-ex].